

Computational Physics


Ising model

- Model
- Numerics
- Examples

Introduction

- The Ising model describes a ferromagnetic (or antiferromagnetic) material.
- A ferromagnet (FM) has a finite magnetization M without applied magnetic field below the Curie temperature T_c .
- At T_c the FM has a second order phase transition to the paramagnetic state at $T > T_c$. M goes to zero at T_c and serves as an order parameter.

The microscopic origin of this macroscopic phenomenon is based on the exchange interaction between identical particles, the atoms or molecules forming the material. The exchange interaction is a ***purely quantum-mechanical effect*** which is a consequence of the ***COULOMB interaction*** in combination with the ***PAULI exclusion principle***.




Spins in a solid – Heisenberg model

Given two atoms or molecules with spins S_1 and S_2 , where $S_1, S_2 \in \mathbb{R}^3$, the exchange interaction energy is of the form:

$$E = JS_1 \cdot S_2$$

with exchange constant J . The ground state for a system of spins depends on the sign of $J \rightarrow$

For the antiferromagnetic case ($J < 0$) the transition temperature T_c is called Neel temperature.



For atoms (spins) on a cubic lattice with lattice points x_ℓ , we can write the Hamilton function

$$H = \frac{1}{2} \sum_{\ell\ell'} J_{\ell\ell'} S_\ell \cdot S_{\ell'} = \frac{1}{2} \sum_{\ell\ell'} J_{\ell-\ell'} S_\ell \cdot S_{\ell'} \quad \text{Heisenberg model}$$

where $J_{\ell\ell} = 0$ and $J_{\ell\ell'} \rightarrow J_{\ell-\ell'} = J_{\ell'-\ell}$ to account for translation invariance.

Ising model

In the Heisenberg model the spin directions are arbitrary. In the Ising model one restricts the direction to one direction, typically the z-direction. Thus the quantum mechanical description (spin quantization) of the (general) Ising model is given by the Hamiltonian

$$H = \frac{1}{2} \sum_{\ell\ell'} J_{\ell-\ell'} S_{\ell}^z S_{\ell'}^z$$

where S_{ℓ}^z are the spin operators in z-direction.

For spin $\frac{1}{2}$ particles, the eigenvalues of the spin operator are $\sigma_{\ell} = \pm 1$ (in units of $\hbar/2$) and we write

$$H = -\frac{1}{2} \sum_{\ell\ell'} J_{\ell-\ell'} \sigma_{\ell} \sigma_{\ell'} - h \sum_{\ell} \sigma_{\ell}$$

**“classical”
Ising model**

where the factor $\hbar^2/4$ is absorbed in the coupling constants and h is an external magnetic field in z-direction.

...

usually the interaction is limited to nearest neighbors (n.n.) such that for $J_{\ell-\ell'}$:

$$J_{\ell-\ell'} = \begin{cases} J & \text{if } \ell, \ell' \text{ n. n.} \\ 0 & \text{otherwise.} \end{cases}$$

$$\rightarrow H = -\frac{J}{2} \sum_{\langle \ell \ell' \rangle} \sigma_{\ell} \sigma_{\ell'} - h \sum_{\ell} \sigma_{\ell}$$



summation over n.n.

For an infinite system, this model can be solved analytically in 1D (Ising's solution) and 2D (Onsager solution)

The magnetization of the system in a certain configuration is given by $\mathcal{M}(C) = \langle \sigma_{\ell} \rangle$.

Statistical Physics and definitions

The Boltzmann distribution is given by: $p(\mathcal{C}) = \frac{1}{Z_N} \exp\left[-\frac{E(\mathcal{C})}{k_B T}\right]$

where $E(\mathcal{C})$ is the energy of a spin configuration \mathcal{C}

The partition function is defined as

$$Z_N = \sum_{\mathcal{C}} \exp\left[-\frac{E(\mathcal{C})}{k_B T}\right]$$

from which we can derive the average energy

$$\langle E \rangle = \sum_{\mathcal{C}} p(\mathcal{C}) E(\mathcal{C}) = k_B T^2 \frac{\partial}{\partial T} \ln Z_N$$

and magnetization

$$\langle M \rangle = \sum_{\mathcal{C}} p(\mathcal{C}) \mathcal{M}(\mathcal{C}) = k_B T \frac{\partial}{\partial h} \ln Z_N$$

using these we can define

$$\chi = \frac{\partial}{\partial h} \langle M \rangle$$

susceptibility

$$c_h = \frac{\partial}{\partial T} \langle E \rangle$$

specific heat

...

Final expressions are:

$$\begin{aligned}c_h &= \frac{1}{k_B} T^2 \sum_{\mathcal{C}} p(\mathcal{C}) [E^2(\mathcal{C}) - E(\mathcal{C}) \langle E \rangle] \\ &= \frac{1}{k_B T^2} (\langle E^2 \rangle - \langle E \rangle^2) \\ &= \frac{1}{k_B T^2} \text{var}(E) .\end{aligned}$$

$$\begin{aligned}\chi &= \frac{1}{k_B T} \sum_{\mathcal{C}} p(\mathcal{C}) [\mathcal{M}^2(\mathcal{C}) - \mathcal{M}(\mathcal{C}) \langle M \rangle] \\ &= \frac{1}{k_B T} (\langle M^2 \rangle - \langle M \rangle^2) \\ &= \frac{1}{k_B T} \text{var}(M) .\end{aligned}$$

1D solution

(see book for details)

The partition function in 1D for N spins can be calculated as

$$Z_N = \lambda_1^N + \lambda_2^N$$

with

$$\lambda_{1,2} = \exp\left(\frac{J}{k_B T}\right) \cosh\left(\frac{h}{k_B T}\right)$$

$$\pm \sqrt{\exp\left(\frac{2J}{k_B T}\right) \sinh^2\left(\frac{h}{k_B T}\right) + \exp\left(-\frac{2J}{k_B T}\right)}$$

The expectation value for the energy per particle is given by

$$\langle \varepsilon \rangle = \frac{k_B T^2}{N} \frac{\partial}{\partial T} \ln Z_N$$

In the thermodynamic limit $N \rightarrow \infty$: $\lim_{N \rightarrow \infty} \frac{1}{N} \ln Z_N = \lim_{N \rightarrow \infty} \frac{1}{N} \ln (\lambda_1^N + \lambda_2^N) = \ln \lambda_1$

and for $h=0$:

$$\lim_{N \rightarrow \infty} \frac{1}{N} \ln Z_N = \ln \left[2 \cosh\left(\frac{J}{k_B T}\right) \right]$$

smooth function of T for $T > 0$
 \rightarrow no phase transition in the
one dimensional Ising model

2D Onsager solution

For $h=0$ one observes a second order phase transition with transition temperature defined by:

$$c_h \text{ and } \chi \text{ diverge at } T_c \quad 2 \tanh^2 \left(\frac{2J}{k_B T_c} \right) = 1 \quad k_B T_c = \frac{2J}{\log(1+\sqrt{2})} \approx 2.269J$$

and for the energy per particle

$$\langle \varepsilon \rangle = -J \coth \left(\frac{2J}{k_B T} \right) \left\{ 1 + \frac{2}{\pi} K_1(\xi) \left[2 \tanh^2 \left(\frac{2J}{k_B T} \right) - 1 \right] \right\}$$

where $K_1(\xi)$ is the complete elliptic integral with

$$\xi = \frac{2 \sinh \left(\frac{2J}{k_B T} \right)}{\cosh^2 \left(\frac{2J}{k_B T} \right)}$$

and magnetization per particle

$$\text{with } z = \exp \left(-\frac{2J}{k_B T} \right)$$

$$\langle m \rangle = \begin{cases} \frac{(1+z^2)^{\frac{1}{4}} (1-6z^2+z^4)^{\frac{1}{8}}}{\sqrt{1-z^2}} & \text{for } T < T_c \\ 0 & \text{for } T > T_c \end{cases}$$

$$= \left(1 - \left[\sinh \left(\log(1 + \sqrt{2}) \frac{T_c}{T} \right) \right]^{-4} \right)^{\frac{1}{8}} \quad T < T_c$$

Phase transitions

Ehrenfest classification of Phase Transition:

- **First-order phase transitions** exhibit a discontinuity in the first derivative of the chemical potential with a thermodynamic variable. Such as solid/liquid/gas transitions.
- **Second-order phase transitions** (also called continuous phase transition) have a discontinuity or divergence in a second derivative of the chemical potential with thermodynamic variables.

c_h and χ are second derivatives

Critical exponents

Reduced temperature: $\tau \equiv \frac{T - T_C}{T_C}$


Critical exponent: $k \stackrel{\text{def}}{=} \lim_{\tau \rightarrow 0} \frac{\log |f(\tau)|}{\log |\tau|}$

Specific heat $C \propto |\tau|^{-\alpha} \quad (T < T_c)$

Magnetization $M \propto |\tau|^\beta$

Magnetic susceptibility $\chi \propto |\tau|^{-\gamma}$

Correlation length $\xi \propto |\tau|^{-\nu}$



Davies, "The New Physics" 1989, p 241

Critical behavior of the order parameter and the correlation length. The order parameter vanishes with the power β of the reduced temperature t as the critical point is approached along the line of phase coexistence. The correlation length diverges with the power ν of the reduced temperature.

The exponents display critical point universality (don't depend on details of the model). This explains the success of the Ising model in providing a quantitative description of real magnets.

Ising values

d	2	3	4
α	0 (log div)	0.110(1)	0
β	1/8	0.3265(3)	1/2
γ	7/4	1.2372(5)	1
δ	15	4.789(2)	3
η	1/4	0.0364(5)	
ν	1	0.6301(4)	1/2
ω	2	0.84(4)	

At $T=T_c$: $H \propto M^\delta$, $\langle \sigma(0)\sigma(r) \rangle \propto r^{2-d-\eta}$

Numerics

Here we concentrate on the 2D case of the Ising model and place the spins on a square lattice Ω with coordinates (x_i, y_j) , $i, j = 1, \dots, N$.

$$H = -J \sum_{\langle ij \rangle} \sigma_{i,j} \sigma_{i',j'} - h \sum_{ij} \sigma_{i,j}$$

with $\sigma_{i,j} = \pm 1$ – i.e. we treat these as “classical” spins. We consider only n.n. interaction.

Task: calculate numerically observables like the expectation value of the energy or of the magnetization, which will then be compared with analytic results.

...

Problem: We cannot sample all possible configurations of the system which grow exponentially with grid points, e.g. for $N=100$ we have $2^{N*N}=2^{10000}\approx 10^{3000}$ possible arrangements of spins.

Solution: Use Monte-Carlo methods – here the Metropolis algorithm

For example, using M (not magnetization here) configurations, we can get an approximation for the energy expectation value

$$\langle E \rangle = \frac{1}{M} \sum_{i=1}^M E(\mathcal{L}_i) \pm \sqrt{\frac{\text{var}(E)}{M}}$$

$$\text{var}(E) = \langle E^2 \rangle - \langle E \rangle^2$$

$$\langle E^2 \rangle = \frac{1}{M} \sum_{i=1}^M E_i^2$$

if these M configurations follow the Boltzmann distribution (this basically means that we neglect most configurations which have exponentially small probability to be physical.)

Metropolis rejection for the Ising model

After generation a new trial configuration \mathcal{C}^t from the current, valid configuration \mathcal{C} , we accept the new one with the following probability

$$\begin{aligned}\Pr(A|\mathcal{C}^t, \mathcal{C}) &= \min\left(\frac{p(\mathcal{C}^t)}{p(\mathcal{C})}, 1\right) = \min\left\{\exp\left[-\frac{E(\mathcal{C}^t) - E(\mathcal{C})}{k_B T}\right], 1\right\} \\ &= \min\left[\exp\left(-\frac{\Delta E_{ij}}{k_B T}\right), 1\right].\end{aligned}$$

\mathcal{C}^t is generated by just flipping one spin on the grid. Since we only consider n.n. interaction, the energy difference of the two configurations is given by the simple expression:

$$\Delta E_{ij} = 2J\sigma_{i,j} (\sigma_{i+1,j} + \sigma_{i-1,j} + \sigma_{i,j-1} + \sigma_{i,j+1}) + 2h\sigma_{i,j}$$

Practical considerations

- We consider a two-dimensional NxN square lattice with periodic boundary conditions in order to reduce finite volume effects

$$\sigma_{N+1,j} = \sigma_{1,j} \quad \text{and} \quad \sigma_{i,N+1} = \sigma_{i,1}$$

- **Do not use** the n.n. matrix suggested in the book – this is usually very inefficient (requires memory access). N.n. coordinates are easy to calculate on-the-fly
- We need a good PRNG to choose random sites
- Initial configuration: the Metropolis algorithm produces configurations which become independent of the initial state and follow the Boltzmann distribution
→ start with random spins
- Run the algorithm for “a while” to **thermalize** the system – disregard these initial trial movements. How to determine “a while”? One should check if thermal equilibrium has been reached by analyzing the observable under consideration as function of time, then determine when initial biases are gone. In this case the observable reaches some saturation value as a function of the number of measurements.

...

- To check if saturation is reached, one should start with two different initial conditions and check when the observables will converge
- typically we study temperature dependencies and change the temperature once the equilibrium has been reached: either hot (random initial spins) or cold (ordered initial spins) start

Running the code

The Metropolis algorithm for the Ising model is executed in the following steps:

1. Choose an initial configuration \mathcal{C}_0
2. we go through all lattice sites (either systematically/sequentially, by random permutation, or completely random) and flip the spin: $\sigma_\ell \rightarrow -\sigma_\ell$. Calculate $\Delta E_{i,j}$ – one complete loop through all site is called a sweep. MC simulations typically need many sweeps
3. Accept the new configuration according to

$$\Pr(A|\mathcal{C}^t, \mathcal{C}_k) = \min \left[\exp \left(-\frac{\Delta E_{ij}}{k_B T} \right), 1 \right]$$



4. go to next lattice site, repeat 2

Measurement:

1. do the above till thermal equilibrium is reached
2. start calculation of observables and average over N trial configurations, when accepted
3. (optional, repeat) change external parameter (T, h), re-equilibrate (typically shorter than the initial equilibration), and then average observable


Some results

$h=0, J=0.5$
 $N=50,$
 $\sim 10^5$ measurements
 (32 sweeps)
 $k_B T = 3 > k_B T_c \rightarrow$
 paramagnetic state,
 $\langle m \rangle = 0$




\rightarrow 32 sweeps are not long enough to equilibrate the energy


temperature dependence



with error bars



domains near T_c



next Homework

- Implement Monte-Carlo solver for the Ising model
 - equilibrate (> 500 sweeps), set $J=0.5$
 - calculate magnetization, average energy, susceptibility and heat capacity across the phase transition
 - do the above to decreasing and increasing temperature for $h=0$
 - calculate error bars
 - choose temperature steps smaller near T_c
 - do the above for $N=5, 20, 50, 100$ or even larger (optional)
 - chose different J and $h \neq 0$ (optional)