

Computational Physics - PHYS 410/510

Spring 2025

Department of Physics - Northern Illinois University
Prof. Andreas Glatz

www.aglatz.net/teaching/compphys_S2025

Homework

1

HW

due 2025-01-28

Info

midterm task: **March 25-27, 2025**
final project presentation: **Thursday, April 24, 2025, 9:30**
(will be assigned beginning of April.)

Program codes should be mailed to: aglatz@niu.edu (see also website). Other problem solutions can be handed in or mailed as well. Problems with points marked by * are for extra credit.

I. COMPUTER ARITHMETIC [2+6+4+6+4 PTS]

- Give the standard (IEEE) single precision binary representation of the machine approximation for $-1/3$.
- Demonstrate how $x_1 + x_2$ and $x_1 - x_2$ are calculated on a computer using decimal floating point numbers with 4 significant digit precision for $x_1 = 0.11258762 \cdot 10^2$ and $x_2 = 0.11244891 \cdot 10^2$. Calculate the relative error.
- Find smallest positive integer that is not exact in single precision.
- How many terms of the exponential series are needed to get the best single and double precision representation of $e = \exp(1)$?
- Calculate the machine epsilon for single and double precision.

II. READ CHAPTER 2.4

III. NUMERICAL DIFFERENTIATION [5+10+15+10+5* PTS]

Consider the finite interval $I = [-a, a]$ with $a = 5$ on the real axis. Define $N + 1$ equally spaced grid-points $x_i = -a + ih$, $i = 0, \dots, N$ spanning I (i.e., $x_N = a$). Investigate the functions

$$g(x) = \exp(-x^2) \text{ and } h(x) = \sin(x),$$

on I .

- Define h . Plot these functions within the interval I by defining these functions on the grid-points x_i for a reasonable N .
- Plot the first derivative of these functions obtained analytically. (use same discretization and N)
- Calculate and plot the first derivatives of these functions by employing the first order backward, forward, and central difference derivatives. (Hand in code). Pay attention to the cases $i = 0$ and $i = N$.
- Find the smallest value of N for each method and function, such that the relative error of numerical and analytical derivative is less than 1%.
- Calculate and plot the second derivative numerically using central differences (see chapter 2.4) for $h(x)$ and find a suitable N such that the relative error is less than 1%.